No.



8300102

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE: PRESENTS SHAYL COME;

Holden's Foundation Seeds, Inc.

Withereas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREONTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE IN THE APPLICANT(S), INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF eighteen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC EED OF THE VARIETY IN A PUBLIC REPOSITORY, AS PROVIDED BY LAW, THE RIGHT TO EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT PORTING IT, OR EXPORTING AT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT 1542, AS AMENDED, 7 U.S.C. 2321 ET SEC.)

CORN

'LH145'

In Lestimony Winercot, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington this 29th day of June in the year of our Lord one thousand nine hundred and eighty-four.

Steet

Kerneth Hoa

Plant Variety Protection Offics Livestock, Meat, Grain & Seed Division Agricultural Marketing Service

Secretary of Agriculture

AGRICULTURAL MAR LIVESTOCK, POULTRY, GR	FORM APPROVED OMB NO. 40-R3822				
- APPEICATION FOR PLANT VARIED INSTRUCTIONS: See Reverse.	No certificate for plant variety protection may be issued unless a completed application form has been received (5 U.S.C. 553).				
1a. TEMPORARY DESIGNATION OF VARIETY	1b. VARIETY NAME		FOR OFFICIAL USE ONLY		
EX 654	LH145		8300 8300	102	
2. KIND NAME	3. GENUS AND SPE	CIES NAME	FILING DATE	TIME A.M.	
	_		4/5/83	9:30 жих.	
Field Corn 4. FAMILY NAME (BOTANICAL)	Zea mays	NAMES TO SECOND	FEE RECEIVED	DATE	
4. PAMILT NAME (BOTANICAL)	5. DATE OF DETER	RMINATION	\$ 1,000 \$ 500.00	4/5/83	
Gramineae	1980		\$ 300.00	5/18/84	
S. NAME OF APPLICANT(S)		and No. or R.F.D. No.,	City, State, and ZIP	8. TELEPHONE AREA	
Had Jan Ja Passa 1-4: au	Code)	0.70		CODE AND NUMBER	
Holden's Foundation . Seeds, Inc.	R.R.#2, I	30x 839 ourg, Iowa 51	2761	319-668-1100	
9. IF THE NAMED APPLICANT IS NOT A P			ED, GIVE STATE AND	11. DATE OF INCOR-	
ORGANIZATION: (Corporation, partners)	nip, association, etc.)	DATE OF INCOR	PORATION	PORATION	
Corporation	·.	Iowa		1968	
12. NAME AND MAILING ADDRESS OF APP ALL PAPERS:		ATIVE(S), IF ANY, TO	SERVE IN THIS APPLIC	CATION AND RECEIVE	
Mr. Art Johns	on				
P.O. Box 839	тана 52361				
Williamsburg 13. CHECK BOX BELOW FOR EACH ATTAC	HMENT SUBMITTED:			,	
🔀 13A. Exhibit A, Origin and Bre	eding History of the	Variety (See Section :	52 of the Plant Variet	y Protection Act.)	
X 13B. Exhibit B, Novelty States	nent.				
13C. Exhibit C, Objective Desc	ription of the Variety	(Request form from	Plant Variety Protect	ion Office.)	
13D. Exhibit D, Additional Des				,	
LAT 13D. Exhibit D, Additional Des	scription of the varies	ry.			
14a. DOES THE APPLICANT(S) SPECIFY THA SEED? (See Section 83(a). (If "Yes," ansu			RIETY NAME ONLY AS	A CLASS OF CERTIFIED	
14b. DOES THE APPLICANT(S) SPECIFY THA LIMITED AS TO NUMBER OF GENERAT			B, HOW MANY GENER BREEDER SEED?	ATIONS OF PRODUC-	
YES X NO		FOUNDATION	REGISTERED	CERTIFIED	
16. DID THE APPLICANT(S) FILE FOR PROT	ECTION OF THIS VAR	RIETY IN OTHER COU	NTRIES? YES	NO (If "Yes," give	
name of countries and dates.)			_		
15b. HAVE RIGHTS BEEN GRANTED THIS V	ARIETY IN OTHER CO	UNTRIES? TYES	NO (If "Yes."	give name of countries	
and dates.)			[X] 140 (1). 123,	give name of countries	
	•				
16. DOES THE APPLICANT(S) AGREE TO THE JOURNAL?	E PUBLICATION OF H	IIS/HER (THEIR) NAM	E(S) AND ADDRESS IN	THE OFFICIAL	
17. The applicant(s) declare(s) that a viab	L	d of this variety will h	e furnished with the	application and will be	
replenished upon request in accordance	e with such regulation	ns as may be applicab	ole.	apparation and will oc	
The undersigned applicant(s) is (are) t	he owner(s) of this se	xually reproduced no	vel plant variety, and	believe(s) that the	
variety is distinct, uniform, and stable 42 of the Plant Variety Act.	as required in Section	n 41, and is entitled t	o protection under th	e provisions of Section	
Applicant(s) is (are) informed that fals	se representation here	in can jeo <u>pard</u> ize pro	tection and result in 1	penalties.	
3/18/93	n de la composition de la composition La composition de la	1/2/1	19/11.		
(DATE)		Monde	SIGNATURE OF APPLI	ICANT)	
	NATIONAL PROPERTY.	gradi.	and the second like		
(DATE)		55	SIGNATURE OF APPLI	ICANT)	

8300102

Page 1

13A Addition Exhibit A:

LH145 was developed through a pedigreed system of breeding. On the following page is a schematic description of the development of LH145. Also included are copies of Holden's Foundation Seeds nursery books. The rows associated with the development of LH145 have been underscored.

Upon observing the increase of LH145 as a finished line for 3 generations, it is free of variance within the population.

Attached is a statement from the originating plant breeder, Art Johnson, Holden's Foundation Seed, stating that the line is uniform and stable.

Uniformity Statement

I observed LH145 during the three generations it has been increased, Iowa Nursery row, 37985, Elwood Garden field 1982 and Hawaii field 1982-83. In each of the increases the seeds from the previous generation were planted. The line is very stable from generation to generation and is very uniform.

Art Johnson Plant Breeder

Exhibit A

Origin and Breeding History of the Variety

$LH145 = Ex654 = A632Ht \times CM105$

Row No.	Pedigree	Location	<u>Year</u>
11562	A632Ht x CM105	Iowa	1976
466	A632Ht x CM105	Iowa	1977
136	A632Ht x CM105	Iowa	1978
5309	A632Ht x CM105	Hawaii	1978-79
7728	A632Ht x CM105	I owa	1979
10109	A632Ht x CM105	Iowa	1980
27985	EX654	Iowa	1981
W. Elwood Gardens	LH145	Iowa	1982
Curtiss: 14-C1	LH145	Hawaii	1982-83

Exhibit B

LH145

LH145 most closely resembles A632Ht; however, the most distinguishing characteristic is the tassel. The lateral branch angle is in the 30° - 40° range for the LH145, while the lateral branch angle of the A632Ht is greater than 45°. This can be seen in the picture on the next page, referred to as Figure 1.




Figure 1.

Addition:

On the right is the LH145 tassel and the A632Ht tassel is on the left.

EXHIBIT C (Corn)

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, POULTRY, GRAIN & SEED DIVISION BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY CORN (ZEA MAYS)

NAME OF APPLICANT(S)	### ##################################
	PYPO NUMBER OF A COMMENT
Holden's Foundation Seeds, Inc. ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	8300102
R.R.#2, Box 839	VARIETY NAME OR TEMPORARY DESIGNATION
Williamsburg, Iowa 52361	
	EX654 or LH145
Place the appropriate number that describes the varietal character of this variety in the Place a zero in first box (e-s-089) or 099) when number is either 99 or less or	
1. TYPE:	
2 1 = SWEET 2 = DENT 3 = FLINT 4 = FLOUR 5 = PC	OP 6 = ORNAMENTAL
2. REGION WHERE BEST ADAPTED IN THE U.S.A.:	
1 = NORTHWEST 2 = NORTHCENTRAL 3 = NORTHEAST 5 = SOUTHCENTRAL 6 = SOUTHWEST 7 = MOST REGIONS	4 = SOUTHEAST
	omments" (pg. 3) state how
	s were calculated)
6, 0 DAYS FROM EMERGENCE TO 50% OF PLANTS IN SILK	9 4 HEAT UNITS
DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY	HEAT UNITS
DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE	HEAT UNITS
4. PLANT:	
2 0 7 CM, HEIGHT (To tassel tip)	8 2 CM, EAR HEIGHT (To base of top ear)
1 5 CM. LENGTH OF TOP EAR INTERNODE	
Number of Tillers: Number of Ears Per Stalk:	
1 1 = NONE 2 = 1-2 3 = 2-3 4 = > 3 2 1 = SINGLE 2	SLIGHT TWO-EAR TENDENCY
3= STRONG TWO	
	EAR TENDENCY 4 = THREE-EAR TENDENCY
Cytoplasm Type:	EAR TENDENCY 4 = THREE-EAR TENDENCY
	•
Cytoplasm Type:	•
Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER	(Specify)Tissues
Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 7.5 Gy 5/6 = Munsell Color charts for Plant	(Specify)Tissues
Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 7.5 Gy 5/6 = Munsell Color charts for Plant 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE Angle from Stalk (Upper half): Sheath Pubscence:	(Specify) Tissues EN (B14) 4 = VERY DARK GREEN (K166)
Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 7.5 Gy 5/6 = Munsell Color charts for Plant 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE Angle from Stalk (Upper half): Sheath Pubscence: 1 1 = LIGHT GREEN (HY) 3 = HEAVY	(Specify) Tissues EN (B14) 4 = VERY DARK GREEN (K166)
Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 7.5 Gy 5/6 = Munsell Color charts for Plant 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° 1 1 = LIGHT (3 = HEAVY)	(Specify) Tissues EN (B14) 4 = VERY DARK GREEN (K166)
Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 7.5 Gy 5/6 = Munsell Color charts for Plant 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = <30° 2 = 30-60° 3 = > 60° 1 = LIGHT GREEN (HE) Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSENT	(Specify) TISSUES EN (B14) 4 = VERY DARK GREEN (K166) W22) 2 = MEDIUM (WF9) (OH26)
Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 7.5 Gy 5/6 = Munsell Color charts for Plant 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° Marginal Waves: Longitudinal Creases:	(Specify) TISSUES EN (B14) 4 = VERY DARK GREEN (K166) W22) 2 = MEDIUM (WF9) (OH26)
Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 7.5 Gy 5/6 = Munsell Color charts for Plant 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° 1 1 = LIGHT (3 = HEAVY) Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSENT 3 = MANY (OH7L)	(Specify) TISSUES EN (B14) 4 = VERY DARK GREEN (K166) W22) 2 = MEDIUM (WF9) (OH26)
Cytoplesm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 7.5 Gy 5/6 = Munsell Color charts for Plant 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° 1 = LIGHT (3 = HEAVY) Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSENT (3 = MANY) (Marginal Creases): Width: Length:	(Specify) TISSUES EN (B14) 4 = VERY DARK GREEN (K166) W22) 2 = MEDIUM (WF9) (OH26)

6. TASSEL:					en e
0 6	受力機を含め、ことをあれた。 NUMBER OF LATERAL		•		
	NUMBER OF LATERAL CANASCAMA AND AND AND AND AND AND AND AND AND AN			· .	
Branch An	le from Central Spike:		Penduncie Length:		
2	1 = < 30. 2 = 30-	-40° 3 × > 45°	0 0 CM. FROM	TOP LEAF TO BASAL BE	IANCHES
	romania de la composición del composición de la	$(\mathbf{z}_{i},\mathbf{y}_{i},\mathbf{y}_{i},\mathbf{y}_{i},\mathbf{y}_{i},\mathbf{y}_{i},\mathbf{y}_{i}) = 0.01$	The second of the second of	, and the second second	
Pollen Shed	. 50		v.		
2	1 = LIGHT (WF9)	2 = MEDIUM	3 = HEAVY(KY21)		
)				
[2.]	,	YELLOW 2 = PINI	к з= RED 4: n_with_purple_stri		GREEN
6	Glume Color: 6 =	OTHER (Specify) greet	i wich barbie zeri	pes	
Pollen Res	oration for Cytoplasms (o	= Not Tested, 1 = Partial, 2 = G	ood)		
<u></u>	 1				
۳۳۰۰ اصا	0 "s"	اما "د" لها ه	THER (Specify Cytoplasm and deg	raes of restoration)	
:					
7. EAR (Hus	ked Ear Data Except When	Stated Otherwise):			
1 6	CM LENGTH 3	MM. MID-POINT DIAMETER	5 8 GM. WEIGH	łT	
Kernel Rov	rs:				
2	1 = INDISTINCT	2 = DISTINCT	1 4 NUMBER		
	1 = STRAIGHT	2 = SLIGHTLY CURVED	3 = SPIRAL		4
Silk Color (Exposed at Silking Stage):				
	1 = GREEN 2	2 = PINK 3 = SALMO	N 4 = RED		
Husk Color	:				
2	FRESH)	I = LIGHT GREEN	2 = DARK GREEN	3 = PINK	
	DRY 4	:= RED . 5= P	URPLE 6 = BUFF	:	
6	tion: (Harvest Stage)		Husk Leaf:		
		MEDIUM (Barely Covering Ear	·	8 CM) 2 = MEDIUM (8-15 CM)
[_2_] 3 = L	DNG (8–10CM Beyond Ea ERY LONG (> 10 CM)	r Tip)	1 3 = LONG (>		
Shank:		•	Position at Dry Husk Stage:		
0 8	CM LONG 7	IO. OF INTERNODES	2 1 = UPRIGHT	2 = HORIZONTAL	3 = PENDENT
Taper:			Drying Time (Unhusked Ear):		
2	1 = SLIGHT 2 = AV	ERAGE 3 = EXTREME	2 1 = SLOW	2 = AVERAGE	3 = FAST
8. KERNEL (I	Oried):				
Size (From	Ear Mid-Point):				
10.	MM LONG	0 5 MM. WIDE) 3 MM. THICK		
Shape Grade	(% Rounds)	· · · · ·			
, 3	1 = < 20 2 = 2	20-40 3 = 40-60	4 = 60 - 80	5 = > 80	

8. KERNE	L (Dried) :							83001	02
8	Pericarp Color:	5 = BROWN		6 = LIGHT R	ite crown ed 1ess brown	7 = CHERI	RY RED	BRONZE	
	Aleurone Color:	1 - HOMOZYG			EGATING (Describe)			o end.	
	1 = WH1TE 7 = PURPLE		3 - 1 JRP LE		4 = BROWN		6 = BR	ONZE	6 = RED
3	Endosperm Color:					4 = PINK	-ORANGE	5 = WHITI	E CAP.
Endospe	rm Type:								
3	1 = SWEET (su1) 5 = WAXY STARC		XTRA SWEE		3 = NORMAL STA 7 = HIGH LYSINE		4 = HIGH AI 8 = OTHER	MYLOSE STAI (Specify)	RCH
2 2	GM, WEIGHT /100	SEEDS (Unsized	Sample)						
9. сов:	MM. DIAMETER A	AT MID-POINT	<u></u>						
Strength	:			Col	or:				
2	1 = WEAK	2 = STRONG		3			= RED 4	l = BROWN ecify)	
10. DISEAS	E RESISTANCE (O	= Not Tested, 1 =	Susceptible, 2	= Resistant):					
0	STALK ROT (Dip!	odia)	0 st	ALK ROT (FI	ışerium)		STALK RO	OT (Gibberella)	i
2	NORTHERN LEAD		o so	UTHERN LEA	AF BLIGHT		SMUT		
	Race 1 southern fust 0 corn smut 0 Bacterial wilt								
0	BACTERIAL LEA	FBLIGHT		NIZE DWARF	MOSAIC		STUNT		
	OTHER (Specify)		<u> </u>		•	الك.ا			
	RESISTANCT (O =	Not Tested, 1 = S	usceptible, 2 =	Resistant):					
	CORNBORER	0	EARWORM		O SAPI	BEETLE		O APHID	
	ROOTWORM (Nor		ROOTWORM						
[0]	ROOTWORM (Sour	thern)	OTHER (Spe	cify)		<u></u>			
		RESEMBLING	THAT SUBMI	TTED FOR T	HE CHARACTERS GI	VEN:			
CHARAC	TER		VARIETY	<u>' </u>	CHARACTER		- V	ARIETY	
Maturity Plant Typ		1 ' '	35Ht32Ht		Kernel Type Quality (Edible	.1	•	632Ht	
Еаг Түрө			32Ht		Usage			 632Ht	
	U.S. Department Ag Corn: Culture, Proc Emerson, R.A., G.W The Mutants of Mais Stringfield, G.H. Ma	essing, Products. Beadle, and A.C 1968. Crop S ize Inbred Lines o	1970 Avi Pub . Fraser. A Sui cience Society of Ohlo, Ohio	mmary of Link of America. A.E.S. Bul. 83		ornell A.E.S.	rous (Authors)	
COMMEN			· · · · · · · · · · · · · · · · · · ·	····			•	1	
	GDD= <u>T</u> n	$\frac{\text{nax} = \text{Tmi}}{2}$	n-50°F		nax = or les nin = or gre			٥ _F	

Exhibit D

LH145

LH145 has some other characteristics that distinguish it from A632Ht. The first is the color. LH145 is lighter in color than A632Ht. When using the Munsell Color Chart for Plant Tissues, the LH145 is classified as 7.5 GY 5/6 and the A632Ht a 7.5 GY 5/4.

Another characteristic is the number of lateral branches. LH145 has fewer lateral branches than A632Ht.

LH145 is shorter than A632Ht in plant height and ear height. The difference is greater for ear height.